



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 5 : A61K 37/02, C07H 21/04, C07K 7/06, 7/08, 15/06, C12N 15/03, 15/06, C12P 21/02, C12Q 1/68	A1	(11) International Publication Number: WO 95/00159 (43) International Publication Date: 5 January 1995 (05.01.95)
(21) International Application Number: PCT/US94/06534 (22) International Filing Date: 10 June 1994 (10.06.94) (30) Priority Data: 08/079,110 17 June 1993 (17.06.93) US 08/196,630 15 February 1994 (15.02.94) US (71) Applicant: LUDWIG INSTITUTE FOR CANCER RE- SEARCH [US/US]; 1345 Avenue of the Americas, New York, NY 10105 (US). (72) Inventors: VAN DER BRUGGEN, Pierre; Avenue Hippocrate 74, UCL 7459, B-1200 Brussels (BE). BOON-FALLEUR, Thierry; Avenue Hippocrate 74, UCL 7459, B-1200 Brussels (BE). COULIE, Pierre; Avenue Hippocrate 74, UCL 7459, B-1200 Brussels (BE). RENAULD, Jean-Christophe; Avenue Hippocrate 74, UCL 7459, B-1200 Brussels (BE). (74) Agent: HANSON, Norman, D.; Felfe & Lynch, 805 3rd Avenue, New York, NY 10022 (US).		(81) Designated States: AU, CA, FI, JP, KR, NO, NZ, European patent (AT, BE, CH, DE, DK, ES, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE). Published <i>With international search report.</i>
(54) Title: ISOLATED PEPTIDES WHICH FORM COMPLEXES WITH MHC MOLECULE HLA-C-CLONE 10 AND USES THEREOF (57) Abstract A new family of tumor rejection antigen precursors, and the nucleic acid molecules which code for them, are disclosed. These tumor rejection antigen precursors are referred to as BAGE tumor rejection antigen precursors, and the nucleic acid molecules which code for them are referred to as BAGE coding molecules. Various diagnostic and therapeutic uses of the coding sequences and the tumorrejection antigen precursor molecules are described.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LI	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

**ISOLATED PEPTIDES WHICH FORM COMPLEXES WITH MHC
MOLECULE HLA-C-CLONE 10 AND USES THEREOF**

RELATED APPLICATION

This application is a continuation-in-part of copending application Serial No. 08/079,110, filed June 17, 1993.

FIELD OF THE INVENTION

This invention relates to peptides which are useful in connection with the diagnosis and treatment of pathological conditions. More particularly, it relates to peptides which are processed to a peptide presented by the MHC molecule HLA-C-clone 10, and the presented peptide itself. These peptides are useful in diagnosis and therapeutic contexts.

BACKGROUND AND PRIOR ART

The process by which the mammalian immune system recognizes and reacts to foreign or alien materials is a complex one. An important facet of the system is the T cell response. This response requires that T cells recognize and interact with complexes of cell surface molecules, referred to as human leukocyte antigens ("HLA"), or major histocompatibility complexes ("MHCs"), and peptides. The peptides are derived from larger molecules which are processed by the cells which also present the HLA/MHC molecule. See in this regard Male et al., Advanced Immunology (J.P. Lipincott Company, 1987), especially chapters 6-10. The interaction of T cell and complexes of HLA/peptide is restricted, requiring a T cell specific for a particular combination of an HLA molecule and a peptide. If a specific T cell is not present, there is no T cell response even if its partner complex is present. Similarly, there is no response if the specific complex is absent, but the T cell is present. This mechanism is involved in the immune system's response to foreign materials, in autoimmune pathologies, and in responses to cellular abnormalities. Much work has focused on the mechanisms by which proteins are processed into the HLA binding peptides. See, in this regard, Barinaga, Science 257: 880 (1992); Fremont et al., Science 257: 919 (1992); Matsumura et al., Science 257: 927 (1992); Latron et al., Science 257:

964 (1992).

The mechanism by which T cells recognize cellular abnormalities has also been implicated in cancer. For example, in PCT application PCT/US92/04354, filed May 22, 1992, published on November 26, 1992, and incorporated by reference, a family of genes is disclosed, which are processed into peptides which, in turn, are expressed on cell surfaces, which can lead to lysis of the tumor cells by specific CTLs. The genes are said to code for "tumor rejection antigen precursors" or "TRAP" molecules, and the peptides derived therefrom are referred to as "tumor rejection antigens" or "TRAs". See Traversari et al., Immunogenetics 35: 145 (1992); van der Bruggen et al., Science 254: 1643 (1991), for further information on this family of genes. Also, see U.S. patent application Serial Number 807,043, filed December 12, 1991, now U.S. Patent No. 5,342,774.

In U.S. patent application Serial Number 938,334, the disclosure of which is incorporated by reference, nonapeptides are taught which are presented by the HLA-A1 molecule. The reference teaches that given the known specificity of particular peptides for particular HLA molecules, one should expect a particular peptide to bind one HLA molecule, but not others. This is important, because different individuals possess different HLA phenotypes. As a result, while identification of a particular peptide as being a partner for a specific HLA molecule has diagnostic and therapeutic ramifications, these are only relevant for individuals with that particular HLA phenotype. There is a need for further work in the area, because cellular abnormalities are not restricted to one particular HLA phenotype, and targeted therapy requires some knowledge of the phenotype of the abnormal cells at issue.

In U.S. Patent Application Serial Number 008,446, filed January 22, 1993 and incorporated by reference, the fact that the MAGE-1 expression product is processed to a second TRA is disclosed. This second TRA is presented by HLA-C clone 10 molecules. The disclosure shows that a given TRAP can yield

5 a plurality of TRAs.

In U.S. Patent Application Serial Number 994,928, filed December 22, 1992, and incorporated by reference herein, tyrosinase is described as a tumor rejection antigen precursor. This reference discloses that a molecule which is
10 produced by some normal cells (e.g., melanocytes), is processed in tumor cells to yield a tumor rejection antigen that is presented by HLA-A2 molecules.

In U.S. patent application Serial No. 08/032,978, filed March 18, 1993, and incorporated by reference in its entirety,
15 a second TRA, not derived from tyrosinase is taught to be presented by HLA-A2 molecules. The TRA is derived from a TRAP, but is coded for by a non MAGE gene. This disclosure shows that a particular HLA molecule may present TRAs derived from different sources.

In U.S. patent application Serial No. 08/079,110 filed June 17, 1993, which is the parent application to the present case and is incorporated by reference in its entirety, a new family of genes, referred to therein as the BAGE family, was disclosed. It was observed that these genes also code for
20 tumor rejection antigen precursors. It was observed in that application that the MHC molecule known as HLA-C-clone 10 presented a tumor rejection antigen derived from a BAGE tumor rejection antigen precursor; however, the tumor rejection antigen was not disclosed. The present application deals with
25 this peptide, referred to herein as SEQ ID NO: 10, as well as other ramifications stemming from the identification of this peptide.

The invention is elaborated upon further in the disclosure which follows.

35 **BRIEF DESCRIPTION OF THE FIGURES**

Figure 1 sets forth the results of a chromium release assay using CTL clone 82/82 against various cell lines.

Figure 2 shows the results of a TNF release assay when CTL clone 82/82 was used against a panel of different cell
40 lines.

5 Figure 3 depicts the results of a TNF release assay using transfected COS cells.

 Figure 4 presents the results of a test of various transfectants using CTL clone 82/82, and measuring TNF release.

10 Figure 5 shows results obtained when studies were undertaken to determine half maximal lysis of cells, following stimulation of cytolytic T cell clone CTL 82/82 with the peptide of SEQ ID NO: 10.

DETAILED DESCRIPTION OF PREFERRED EMBODIMENTS

15 Example 1

 A melanoma cell line, MZ2-MEL was established from melanoma cells taken from patient MZ2, using standard methodologies. This cell line is described, e.g., in PCT Application PCT/US92/04354, filed May 22, 1992, published
20 November 26, 1992, and incorporated by reference in its entirety. Once the cell line was established, a sample thereof was irradiated, so as to render it non-proliferative. These irradiated cells were then used to isolate cytolytic T cell clones ("CTLs") specific thereto.

25 A sample of peripheral blood mononuclear cells ("PBMCs") was taken from patient MZ2, and contacted to the irradiated melanoma cells. The mixture was observed for lysis of the melanoma cells, which indicated that CTLs specific for a complex of peptide and HLA molecule presented by the melanoma
30 cells were present in the sample.

 The lysis assay employed was a chromium release assay following Herin et al., Int. J. Cancer 39:390-396 (1987), the disclosure of which is incorporated by reference. The assay, however, is described herein. The target melanoma cells were
35 grown in vitro, and then resuspended at 10^7 cells/ml in DMEM, supplemented with 10 mM HEPES and 30% FCS, and incubated for 45 minutes at 37°C with 200 μ Ci/ml of $\text{Na}^{51}\text{Cr}\text{O}_4$. Labelled cells were washed three times with DMEM, supplemented with 10 mM Hepes. These were then resuspended in DMEM supplemented
40 with 10 mM Hepes and 10% FCS, after which 100 μ l aliquots containing 10^3 cells, were distributed into 96 well

5 microplates. Samples of PBLs were added in 100 ul of the same medium, and assays were carried out in duplicate. Plates were centrifuged for 4 minutes at 100g, and incubated for four hours at 37°C in a 8% CO₂ atmosphere.

Plates were centrifuged again, and 100 ul aliquots of
10 supernatant were collected and counted. Percentage of ⁵¹Cr release was calculated as follows:

$$\% \text{ } ^{51}\text{Cr release} = \frac{(\text{ER}-\text{SR})}{(\text{MR}-\text{SR})} \times 100$$

15

where ER is observed, experimental ⁵¹Cr release, SR is spontaneous release measured by incubating 10³ labeled cells in 200 ul of medium alone, and MR is maximum release, obtained by adding 100 ul 0.3% Triton X-100 to target cells.

20 Those mononuclear blood samples which showed high CTL activity were expanded and cloned via limiting dilution, and were screened again, using the same methodology. The CTL clone MZ2-CTL 82/82 was thus isolated. The clone is referred to as "82/82" hereafter.

25 The same method was used to test target K562 cells, as well as a melanoma cell line. These results, presented in Figure 1A, show that this CTL clone recognizes and lyses the melanoma cell line, but not K562. The CTL clone, 82/82, was then tested against melanoma cell lines in the same manner described supra. Figure 1 shows that, while MZ2-MEL.43 is
30 lysed by CTL clone 82/82, the cell line MZ2-MEL 2.2.5, a variant which has lost expression of HLA-A29, HLA-B44, and HLA-C clone 10 is not, suggesting that the TRA is presented by one of these HLA molecules. When cell line MZ2-MEL 2.2.5 was
35 transfected with DNA coding for HLA-C clone 10, using well known techniques, the cells became sensitive to lysis by CTL clone 82/82, thus demonstrating that the antigen recognized by CTL clone 82/82 is presented by HLA-C clone 10.

5 Example 2

Further studies were carried out to determine if 82/82 also produced tumor necrosis factor ("TNF") when contacted with target cells. The method used was that described by Traversari et al., Immunogenetics 35: 145-152 (1992), the disclosure of which is incorporated by reference. Briefly, 10 samples of the CTL line were combined with samples of a target cell of interest, in culture medium. After 24 hours, supernatant from the cultures was removed, and then tested on TNF sensitive WEHI cells. A panel of fourteen different 15 cell lines were tested, as shown in Figure 2.

The results, presented in terms of the percentage of WEHI cells which died upon exposure to the supernatant, are shown in Figure 2. The results demonstrate that three melanoma cell lines present this antigen. As a result of the strong 20 response by MZ2 MEL43, it was used in the experiments which follow.

Example 3

The results from Example 2 indicated that MZ2.MEL.43 25 presented the target antigen of interest. As such, it was used as a source of total mRNA to prepare a cDNA library.

Total RNA was isolated from the cell line. The mRNA was isolated using an oligo-dT binding kit, following well recognized techniques. Once the mRNA was secured, it was 30 transcribed into cDNA, again using standard methodologies. The cDNA was then ligated to EcoRI adaptors and cloned into the EcoRI site of plasmid pcD-SR α , in accordance with the manufacturer's instructions. The recombinant plasmids were then electroporated into JM101 *E. coli* (electroporation 35 conditions: 1 pulse at 25 μ farads, 2500 V).

The transfected bacteria were selected with ampicillin (50 μ g/ml), and then divided into 87 pools of 400 bacteria and 297 pools of 200 bacteria. Each pool represented either about 280 or about 140 cDNAs, as analysis showed that about 40 70% of plasmids contained an insert. Each pool was amplified to saturation, and plasmid DNA was isolated via alkaline

5 lysis, potassium acetate precipitation without phenol extraction, following Maniatis et al., in Molecular Cloning: A Laboratory Manual (Cold Spring Harbor, N.Y., 1982).

Example 4

10 Following preparation of the library described in Example 3, the cDNA was transfected into eukaryotic cells. The transfections, described herein, were carried out in duplicate. Samples of COS-7 cells were seeded, at 15,000 cells/well into tissue culture flat bottom microwells, in Dulbecco's modified Eagles Medium ("DMEM") supplemented with
15 10% fetal calf serum. The cells were incubated overnight at 37°C, medium was removed and then replaced by 50 µl/well of DMEM medium containing 10% Nu serum, 400 µg/ml DEAE-dextran, and 100 µM chloroquine, plus 100 ng of subject plasmids. These plasmids were the plasmids of the various pools
20 described supra, and 100 ng of plasmids containing DNA coding for HLA-C clone 10 in plasmid pcD-SRα. Following four hours of incubation at 37°C, the medium was removed, and replaced by 50 µl of PBS containing 10% DMSO. This medium was removed after two minutes and replaced by 200 µl of DMEM supplemented
25 with 10% FCS.

Following this change in medium, COS cells were incubated for 24-48 hours at 37°C. Medium was then discarded, and 1500 cells of CTL clone 82/82 were added, in 100 µl of Iscove medium containing 10% pooled human serum supplemented with
30 U/ml of IL-2. Supernatant was removed after 24 hours, and TNF content was determined in an assay on WEHI cells, as described by Traversari et al., Immunogenetics 35: 145-152 (1992), the disclosure of which is incorporated by reference.

35 Of the 384 pools tested, 99% stimulated TNF, at a concentration below 5 pg/ml. Two pools gave concentrations above 40 pg/ml, with duplicate wells giving equivalent results. Figure 3 shows this. The bacteria from one of these pools, i.e., pool 19, were selected for further experiments.

5 Example 5

 The bacteria of pool 19 were cloned, and 800 bacteria were tested. Plasmid DNA was extracted therefrom, transfected into a new sample of COS cells in the same manner as described supra, and the cells were again tested for stimulation of CTL clone 82/82. Twelve positive clones were found. One of these, referred to as cDNA clone AD5 was tested further. In a comparative test COS cells were transfected with cDNA clone AD5 and the HLA-C clone 10. HLA-C clone 10 and MAGE-1, AD5 alone, or HLA-C clone 10 alone. Control cell lines MZ2-MEL 2.2.5 and MZ2-MEL.43 were also used. TNF release in CTL supernatant was measured by testing it on WEHI cells, as referred to supra. The optical density of the surviving WEHI cells was measured using MTT. Figure 4 shows that only the COS cells transfected with HLA-C clone 10 and cDNA-AD5, and the original cell line MZ2-MEL 43 stimulated TNF release from CTL clone 82/82.

Example 6

 The cDNA AD5 was sequenced following art known techniques. A sequence search revealed that the plasmid insert showed no homology to known genes or proteins. SEQUENCE ID NO: 1 presents cDNA nucleotide information for the identified gene, referred to hereafter as "BAGE-1". A putative open reading frame is located at bases 201-332 of the molecule.

30 Example 7

 Following the sequencing of the cDNA, as per example 6, experiments were carried out to determine if cells of normal tissues expressed the gene. To determine this, RNA isolated from normal tissues was reverse transcribed, using oligo-dT as primer. The resulting cDNA was then amplified, using primers:

 5' CAG AAG ATG AAG CAC AGA G-3' (SEQ ID NO: 2)

and

 5'-GAG CGG TTT TTC TGG CAT TG-3' (SEQ ID NO: 3)

and standard PCR methodologies. Radioactive nucleotides were

5 added so that the amount of amplification product could be determined via phosphor imaging.

The amount of product was expressed as a percentage of the product secured from cell line MZ2-MEL 3.0, which was shown to express the gene. The results are as follows:

10	MZ2-MEL 3.0	100%
	Lung	< 0.5%
	Breast	"
	Stomach	"
	Skin	"
15	Brain	"
	Prostate	"
	Kidney	"
	Testis	8%

20 In additional experiments not elaborated upon herein, the DNA of cell line MZ2-MEL was digested with EcoRI, and then hybridized with a PCR probe corresponding to the first 300 nucleotides of the cDNA described herein. Following standard Southern blotting, four bands, corresponding to sizes of approximately 5.8, 7.5, 8.5 and 11 kilobases were identified, suggesting the existence of a family of Bage genes.

25 Example 8

Expression of the gene by tumor samples and tumor cell lines was also determined. cDNA was secured just as in example 4, and then nested primer methodologies were carried out to amplify the pertinent sequences. First, twenty cycles of amplification were carried out, using primers:

5'-CGG CTT AGA GGA CCA GGA GAA-3' (SEQ ID NO: 4)

and

5'CAG AAG ATG AAG CAC AGA G-3' (SEQ ID NO: 5)

35

This was followed by twenty additional cycles using primers:

5'-GGC TCC AAC CTC CAG CTC AC-3' (SEQ ID NO: 6)

AND

5'-TTA GAG GAC CAG GAG AAG G-3' (SEQ ID NO: 7)

40

The results are presented below. The first figure is the

10

5	number of positiv samples; the second is the total number of samples tested:	
	Melanoma	12/20
	Breast Carcinoma	2/5
	Small Cell lung carcinoma	2/8
10	Non-small cell lung carcinoma	2/5
	Sarcoma	1/4
	Head and neck tumors	1/6
	Colon carcinoma	0/4
	Kidney tumor	0/5
15	Leukemia/Lymphoma	0/3

Example 9

The experiments set forth supra showed that cytolytic T cell clone CTL 82/82 recognized an antigen encoded by a BAGE gene and presented by HLA-C-clone 10. The work described in this example details how the amino acid sequence of the presented antigen was determined.

The cDNA clone identified as coding for BAGE, i.e., AD5, was used to generate a large number of incomplete cDNA molecules. The cDNA clone was inserted into expression vector pcDNAI/Amp, and digested with NotI and SphI restriction endonucleases, followed by treatment with exonuclease III, in accordance with the manufacturer's instructions. By using exonuclease III for varying lengths of time, progressive deletions of AD5 at its 3' end were obtained. The truncated variants were religated into pcDNAI/Amp, electroporated into *E. coli* strain DH5 α F'IQ, and selected via ampicillin (50 μ g/ml). Four hundred clones were obtained in this way.

The plasmid DNA was obtained from these 400 clones, and transfected into COS-7 cells together with HLA-C-clone 10 coding cDNA. The transfectants were then tested in a TNF release assay, as described supra. Positive clones were those which stimulated TNF release by CTL 82/82.

Once cells were divided into positive and negative transfectants, the sequences of plasmid DNA from 10 positives and 10 negatives was determined. Clone 19C2, a positive

clone, contained part of the open reading frame for the BAGE gene described supra, from nucleotide 1 to nucleotide 67. In contrast, clone 17G12, a negative transfectant, contained nucleotides 1-6 of the gene.

By comparing the inserts of positive and negative clones, a region of 22 amino acids was identified as probably containing the sequence of the presented peptide, i.e.:

Met Ala Ala Arg Ala Val Phe Leu Ala Leu Ser Ala Gln Leu Leu Gln
Ala Arg Leu Met Lys Glu
(SEQ ID NO: 8)

Synthetic peptides were then prepared, based upon this sequence, and were tested for their ability to render COS-7 cells transfected with HLA-C-clone 10 capable of stimulating TNF release. The first positive peptide was a 16-mer:

Met Ala Ala Arg Ala Val Phe Leu Ala Leu Ser Ala Gln Leu Leu Gln
(SEQ ID NO: 9)

Testing of smaller peptides led to the identification of peptide

Ala Ala Arg Ala Val Phe Leu Ala Leu
(SEQ ID NO: 10)

This peptide effectively stimulated CTL 82/82, with half maximal lysis being reached at a peptide concentration of 80 nM, which is shown in Figure 5.

The foregoing examples show the isolation of a nucleic acid molecule which codes for a tumor rejection antigen precursor. This "TRAP" coding molecule, however, is not homologous with any of the previously disclosed MAGE coding sequences described in the references set forth supra. Hence, one aspect of the invention is an isolated nucleic acid molecule which comprises the nucleotide sequence set forth in SEQ ID NO: 1. This sequence is not a MAGE coding sequence, as will be seen by comparing it to the sequence of any of the MAGE genes described in the references. Also a part of the invention are those nucleic acid sequences which also code for a non-MAGE tumor rejection antigen precursor but which hybridize to a nucleic acid molecule containing the described nucleotide sequence, under stringent conditions. The term "stringent conditions" as used herein refers to parameters

with which the art is familiar. More specifically, stringent conditions, as used herein, refers to hybridization in 3.5xSSC, 1xDenhardt's solution, 25 mM sodium phosphate buffer (pH 7.0), 0.5% SDS, and 2 mM EDTA for 18 hours at 65°C. This is followed by four washes of the filter, at 65°C for 20 minutes, in 2xSSC, 0.1% SDS, and one wash for up to 20 minutes in 0.3xSSC, 0.1% SDS. There are other conditions, reagents, and so forth which can be used, which result in the same degree of stringency. The skilled artisan will be familiar with such conditions, and thus they are not given here.

It will also be seen from the examples that the invention embraces the use of the sequences in expression vectors, as well as to transfect host cells and cell lines, be these prokaryotic (e.g., *E. coli*), or eukaryotic (e.g., CHO or COS cells). The expression vectors require that the pertinent sequence, i.e., those described supra, be operably linked to a promoter. As it has been found that human leukocyte antigen HLA-C clone 10 presents a tumor rejection antigen derived from these genes, the expression vector may also include a nucleic acid sequence coding for HLA-C clone 10. In a situation where the vector contains both coding sequences, it can be used to transfect a cell which does not normally express either one. The tumor rejection antigen precursor coding sequence may be used alone, when, e.g., the host cell already expresses HLA-C clone 10. Of course, there is no limit on the particular host cell which can be used. As the vectors which contain the two coding sequences may be used in HLA-C clone 10 presenting cells if desired, and the gene for tumor rejection antigen precursor can be used in host cells which do not express HLA-C clone 10.

The invention also embraces so called expression kits, which allow the artisan to prepare a desired expression vector or vectors. Such expression kits include at least separate portions of each of the previously discussed coding sequences. Other components may be added, as desired, as long as the previously mentioned sequences, which are required, are included.

To distinguish the nucleic acid molecules and the TRAPs of the invention from the previously described MAGE family, the invention shall be referred to as the BAGE family of genes and TRAPs. Hence, whenever "BAGE" is used herein, it refers to the tumor rejection antigen precursors coded for by the previously described sequences. "BAGE coding molecule" and similar terms, are used to describe the nucleic acid molecules themselves.

Also a part of the invention are the peptides of SEQ ID NO: 8, 9 and 10 which are set forth in Example 9. These peptides can be used, for example, to identify those cells which present MHC molecule HLA-C-clone 10. Administration of the peptides, carrying a detectable signal, e.g., followed by the identification of cells to which the peptide has bound is one way to accomplish this, as is the use of solid phase bound peptides, to which the HLA-C-clone 10 presenting cells bind, thus removing them from the sample being assayed. Additionally, the invention permits the artisan to diagnose a disorder characterized by expression of the TRAP. These methods involve determining expression of the TRAP gene, and/or TRAs derived therefrom, such as the TRA presented by HLA-C clone 10. In the former situation, such determinations can be carried out via any standard nucleic acid determination assay, including the polymerase chain reaction, or assaying with labelled hybridization probes. In the latter situation, assaying with binding partners for complexes of TRA and HLA, such as antibodies, is especially preferred. An alternate method for determination is a TNF release assay, of the type described supra.

The isolation of the TRAP gene also makes it possible to isolate the TRAP molecule itself, especially TRAP molecules containing the amino acid sequence coded for by SEQ ID NO: 1. These isolated molecules when presented as the TRA, or as complexes of TRA and HLA, such as HLA-C clone 10, may be combined with materials such as adjuvants to produce vaccines useful in treating disorders characterized by expression of the TRAP molecule. In addition, vaccines can be prepared from

cells which present the TRA/HLA complexes on their surface, such as non-proliferative cancer cells, non-proliferative transfectants, etcetera. In all cases where cells are used as a vaccine, these can be cells transfected with coding sequences for one or both of the components necessary to prove a CTL response, or be cells which express both molecules without transfection. Further, the TRAP molecule, its associated TRAs, as well as complexes of TRA and HLA, may be used to produce antibodies, using standard techniques well known to the art.

When "disorder" is used herein, it refers to any pathological condition where the tumor rejection antigen precursor is expressed. An example of such a disorder is cancer melanoma in particular.

Therapeutic approaches based upon the disclosure are premised on a response by a subject's immune system, leading to lysis of TRA presenting cells, such as HLA-c clone 10 cells. One such approach is the administration of CTLs specific to the complex to a subject with abnormal cells of the phenotype at issue. It is within the skill of the artisan to develop such CTLs in vitro. Specifically, a sample of cells, such as blood cells, are contacted to a cell presenting the complex and capable of provoking a specific CTL to proliferate. The target cell can be a transfectant, such as a COS cell of the type described supra. These transfectants present the desired complex on their surface and, when combined with a CTL of interest, stimulate its proliferation.

COS cells, such as those used herein are widely available, as are other suitable host cells.

To detail the therapeutic methodology, referred to as adoptive transfer (Greenberg, J. Immunol. 136(5): 1917 (1986); Reddel et al., Science 257: 238 (7-10-92); Lynch et al., Eur. J. Immunol. 21: 1403-1410 (1991); Kast et al., Cell 59: 603-614 (11-17-89)), cells presenting the desired complex are combined with CTLs leading to proliferation of the CTLs specific thereto. The proliferated CTLs are then administered to a subject with a cellular abnormality which is

characterized by certain of the abnormal cells presenting the particular complex. The CTLs then lyse the abnormal cells, thereby achieving the desired therapeutic goal.

The foregoing therapy assumes that at least some of the subject's abnormal cells present the relevant HLA/TRA complex. This can be determined very easily, as the art is very familiar with methods for identifying cells which present a particular HLA molecule, as well as how to identify cells expressing DNA of the pertinent sequences, in this case a BAGE sequence. Once cells presenting the relevant complex are identified via the foregoing screening methodology, they can be combined with a sample from a patient, where the sample contains CTLs. If the complex presenting cells are lysed by the mixed CTL sample, then it can be assumed that a BAGE derived, tumor rejection antigen is being presented, and the subject is an appropriate candidate for the therapeutic approaches set forth supra.

Adoptive transfer is not the only form of therapy that is available in accordance with the invention. CTLs can also be provoked in vivo, using a number of approaches. One approach, i.e., the use of non-proliferative cells expressing the complex, has been elaborated upon supra. The cells used in this approach may be those that normally express the complex, such as irradiated melanoma cells or cells transfected with one or both of the genes necessary for presentation of the complex. Chen et al., Proc. Natl. Acad. Sci. USA 88: 110-114 (January, 1991) exemplifies this approach, showing the use of transfected cells expressing HPVE7 peptides in a therapeutic regime. Various cell types may be used. Similarly, vectors carrying one or both of the genes of interest may be used. Viral or bacterial vectors are especially preferred. In these systems, the gene of interest is carried by, e.g., a Vaccinia virus or the bacteria BCG, and the materials de facto "infect" host cells. The cells which result present the complex of interest, and are recognized by autologous CTLs, which then proliferate. A similar effect can be achieved by combining the tumor rejection antigen or the precursor itself with an

adjuvant to facilitate incorporation into HLA-C clone 10 presenting cells which present the HLA molecule of interest. The TRAP is processed to yield the peptide partner of the HLA molecule while the TRA is presented without the need for further processing.

Other aspects of the invention will be clear to the skilled artisan and need not be repeated here.

The terms and expressions which have been employed are used as terms of description and not of limitation, and there is no intention in the use of such terms and expressions of excluding any equivalents of the features shown and described or portions thereof, it being recognized that various modifications are possible within the scope of the invention.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: van der Bruggen, Pierre
Boon-Falleur, Thierry
- (ii) TITLE OF INVENTION: ISOLATED PEPTIDES WHICH FORM COMPLEXES WITH
MHC MOLECULE HLA-C-CLONE 10 AND USES THEREOF
- (iii) NUMBER OF SEQUENCES: 10
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Felfe & Lynch
 - (B) STREET: 805 Third Avenue
 - (C) CITY: New York City
 - (D) STATE: New York
 - (E) COUNTRY: USA
 - (F) ZIP: 10022
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Diskette, 5.25 inch, 360 kb storage
 - (B) COMPUTER: IBM PS/2
 - (C) OPERATING SYSTEM: PC-DOS
 - (D) SOFTWARE: Wordperfect
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/196,630
 - (B) FILING DATE: 15-FEB-1994
 - (C) CLASSIFICATION: 435
- (vii) PRIOR APPLICATION DATA:
 - (A) APPLICATION NUMBER: 08/079,110
 - (B) FILING DATE: 17-JUN-1993
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Hanson, Norman D.
 - (B) REGISTRATION NUMBER: 30,946
 - (C) REFERENCE/DOCKET NUMBER: LUD 310.1
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: (212) 688-9200
 - (B) TELEFAX: (212) 838-3884

18

- (2) INFORMATION FOR SEQ ID NO: 1:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 1032 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 1:

```
CGCCAATTTA GGGTCTCCGG TATCTCCCGC TGAGCTGCTC TGTTCCCGGC TTAGAGGACC 60
AGGAGAAGGG GGAGCTGGAG GCTGGAGCCT GTAACACCGT GGCTCGTCTC ACTCTGGATG 120
GTGGTGGCAA CAGAGATGGC AGCGCAGCTG GAGTGTTAGG AGGGCGGCCT GAGCGGTAGG 180
AGTGGGGCTG GAGCAGTAAG ATGGCGGCCA GAGCGGTTT TCTGGCATTG TCTGCCCAGC 240
TGCTCCAAGC CAGGCTGATG AAGGAGGAGT CCCCTGTGGT GAGCTGGAGG TTGGAGCCTG 300
AAGACGGCAC AGCTCTGTGC TTCATCTTCT GAGGTTGTGG CAGCCACGGT GATGGAGACG 360
GCAGCTCAAC AGGAGCAATA GGAGGAGATG GAGTTTCACT GTGTCAGCCA GGATGGTCTC 420
GATCTCCTGA CCTCGTGATC CGCCCGCCTT GGCCTTCCAA AGTGCCGAGA TTACAGCGAT 480
GTGCATTTTG TAAGCACTTT GGAGCCACTA TCAAATGCTG TGAAGAGAAA TGTACCCAGA 540
TGTATCATTG TCCTTGTGCT GCAGGAGCCG GCTCCTTTCA GGATTTTCAGT CACATCTTCC 600
TGCTTTGTCC AGAACACATT GACCAAGCTC CTGAAAGATG TAAGTTTACT ACGCATAGAC 660
TTTTAACTT CAACCAATGT ATTTACTGAA AATAACAAAT GTTGTAATTT CCCTGAGTGT 720
TATTCTACTT GTATTAAAAG GTAATAATAC ATAATCATTG AAATCTGAGG GATCATTGCC 780
AGAGATTGTT GGGGAGGGAA ATGTTATCAA CGGTTTCATT GAAATTAAAT GTTATCAACG 840
GTTTCATTGA AATTAAATCC AAAAAATTAT TTCCTCAGAA AAATCAAATA AAGTTTGCAT 900
GTTTTTTTATT CTTAAAACAT TTTAAAACC ACTGTAGAAT GATGTAAATA GGGACTGTGC 960
AGTATTTCTG ACATATACTA TAAATTATT AAAAAGTCAA TCAGTATTCA ACATCTTTTA 1020
CACTAAAAAG CC 1032
```

- (2) INFORMATION FOR SEQ ID NO: 2:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 2:

CAGAAGATGA AGCACAGAG

9

- (2) INFORMATION FOR SEQ ID NO: 3:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 3:

GAGCGGTTTT TCTGGCATTG

20

- (2) INFORMATION FOR SEQ ID NO: 4:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 21 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 4:

19

CGGCTTAGAG GACCAGGAGA A

21

- (2) INFORMATION FOR SEQ ID NO: 5:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 5:

CAGAAGATGA AGCACAGAG

19

- (2) INFORMATION FOR SEQ ID NO: 6:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 20 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 6:

GGCTCCAACC TCCAGCTCAC

20

- (2) INFORMATION FOR SEQ ID NO: 7:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 19 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 7:

TTAGAGGACC AGGAGAAGG

19

- (2) INFORMATION FOR SEQ ID NO: 8:
 (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 22 amino acid residues
 (B) TYPE: amino acid
 (D) TOPOLOGY: single
 (xi) SEQUENCE DESCRIPTION: SEQ ID NO: 8:

Met Ala Ala Arg Ala Val Phe Leu Ala Leu Ser Ala Gln Leu Leu Gln
 5 10 15
 Ala Arg Leu Met Lys Glu
 20

- (2) INFORMATION FOR SEQ ID NO: 9:
 (i) SEQUENCE CHARACTERISTICS:

20

(A) LENGTH: 16 amino acid residues
(B) TYPE: amino acid
(D) TOPOLOGY: single
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 9:

Met Ala Ala Arg Ala Val Phe Leu Ala Leu Ser Ala Gln Leu Leu Gln
5 10 15

(2) INFORMATION FOR SEQ ID NO: 10:

(i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 9 amino acid residues
(B) TYPE: amino acid
(D) TOPOLOGY: single
(xi) SEQUENCE DESCRIPTION: SEQ ID NO: 10:

Ala Ala Arg Ala Val Phe Leu Ala Leu
5

21

W claim:

1. An isolated nucleic acid molecule comprising the nucleotide sequence set forth in SEQ ID NO: 1.
2. An isolated nucleic acid molecule which hybridizes, under stringent conditions, to the nucleic acid sequence set forth in SEQ ID NO: 1, and codes for a tumor rejection antigen precursor, with the proviso that said isolated nucleic acid molecule does not code for a MAGE tumor rejection antigen precursor.
3. An isolated mRNA molecule which is complementary to the nucleic acid molecule of claim 1.
4. A host cell transfected with the nucleic acid molecule of claim 1.
5. A host cell transfected with the nucleic acid molecule of claim 2.
6. An expression vector comprising the isolated nucleic acid molecule of claim 1 operably linked to a promoter.
7. An expression vector comprising the isolated nucleic acid molecule of claim 2 operably linked to a promoter.
8. The host cell of claim 4, wherein said host cell is a mammalian cell which expresses HLA-C clone 10.
9. The host cell of claim 5, wherein said host cell is a mammalian cell which expresses HLA-C clone 10.
10. The expression vector of claim 6, further comprising a nucleic acid molecule which codes for HLA-C clone 10.
11. The expression vector of claim 7, further comprises a nucleic acid molecule which codes for HLA-C clone 10.
12. Expression kit comprising a separate portion of each of:
 - (i) the isolated nucleic acid molecule of claim 1, and
 - (ii) a nucleic acid molecule which codes for HLA-C clone 10.
13. Expression kit comprising a separate portion of each of:
 - (i) the isolated nucleic acid molecule of claim 2, and;
 - (ii) a nucleic acid molecule which codes for HLA-C clone 10.
14. An isolated tumor rejection antigen precursor coded for by the nucleic acid molecule of claim 1.

15. Method for treating a subject with a disorder characterized by expression of a BAGE tumor rejection antigen precursor which is processed to a tumor rejection antigen presented by HLA-C clone 10 molecules comprising: administering to said subject an amount of cytolytic T cells which are specific for complexes of said BAGE derived tumor rejection antigen and HLA-C clone 10 molecules and which lyse cells presenting said complexes, sufficient to alleviate said disorder.
16. Method for treating a subject with a disorder characterized by expression of a tumor rejection antigen precursor coded for by a nucleic acid molecule and comprising the nucleotide sequence of SEQ ID NO: 1, comprising administering to said subject an amount of cytolytic T cells specific to complexes of an HLA molecule and a tumor rejection antigen derived from said tumor rejection antigen precursor, sufficient to alleviate said disorder.
17. Method for treating a subject with a disorder characterized by expression of a BAGE tumor rejection antigen precursor which is processed to a tumor rejection antigen presented by HLA-C clone 10 molecules, comprising administering to said subject an amount of an agent which provokes an immune response to complexes of said BAGE derived tumor rejection antigen and HLA-C clone 10 molecules sufficient to provoke said response against cells presenting said complex.
18. Method for treating a subject with a disorder characterized by expression of a tumor rejection antigen precursor coded for by a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 1, comprising administering to said subject an amount of an agent which provokes an immune response to complexes of an HLA molecule and a tumor rejection antigen precursor, sufficient to provoke said immune response against cells presenting said complexes.
19. Method for diagnosing a disorder characterized by

expression of a BAGE tumor rejection antigen precursor which is processed to a BAGE derived tumor rejection antigen which forms a complex with HLA-C clone 10 molecules, comprising contacting a sample from a subject with an agent specific for said complex and determining interaction between said complex and said agent as a determination of said disorder.

20. Method for diagnosing a disorder characterized by expression of a tumor rejection antigen precursor coded for by a nucleic acid molecule having the sequence set forth in SEQ ID NO: 1, comprising contacting a sample from a subject with an antigen specific for said sequence or an expression product thereof, and determining interaction between said agent and said sequence or said expression product as a determination of said disorder.
21. Isolated peptide selected from the group consisting of: SEQ ID NO: 8, SEQ ID NO: 9, and SEQ ID NO: 10.
22. Method for provoking a cytolytic T cell response comprising contacting an HLA-C-clone 10 presenting cell with the isolated peptide of claim 21 in the presence of cytolytic T cells, in an amount sufficient to provoke proliferation of cytolytic T cells specific to complexes of HLA-C-clone 10 and SEQ ID NO: 10.
23. Method for treating a subject with a disorder characterized by expression of a BAGE tumor rejection antigen precursor which is processed to a tumor rejection antigen consisting of the amino acid sequence of SEQ ID NO: 10 which is presented by HLA-C clone 10 molecules comprising: administering to said subject an amount of cytolytic T cells which are specific for complexes of said BAGE derived tumor rejection antigen and HLA-C clone 10 molecules and which lyse cells presenting said complexes, sufficient to alleviate said disorder.
24. Method for treating a subject with a disorder characterized by expression of a tumor rejection antigen precursor coded for by a nucleic acid molecule and comprising the nucleotide sequence of SEQ ID NO: 1,

comprising administering to said subject an amount of cytolytic T cells specific to complexes of an HLA molecule and a tumor rejection antigen consisting of the amino acid sequence of SEQ ID NO: 10, sufficient to alleviate said disorder.

25. Method for treating a subject with a disorder characterized by expression of a BAGE tumor rejection antigen precursor which is processed to a tumor rejection antigen consisting of the amino acid sequence of SEQ ID NO: 10 which is presented by HLA-C clone 10 molecules, comprising administering to said subject an amount of an agent which provokes an immune response to complexes of said BAGE derived tumor rejection antigen and HLA-C clone 10 molecules sufficient to provoke said response against cells presenting said complex.
26. Method for treating a subject with a disorder characterized by expression of a tumor rejection antigen precursor coded for by a nucleic acid molecule comprising the nucleotide sequence of SEQ ID NO: 1, comprising administering to said subject an amount of an agent which provokes an immune response to complexes of HLA-C-clone and a peptide consisting of the amino acid sequence of SEQ ID NO: 10, sufficient to provoke said immune response against cells presenting said complexes.
27. Method for diagnosing a disorder characterized by expression of a BAGE tumor rejection antigen precursor which is processed to a BAGE derived tumor rejection antigen consisting of the amino acid sequence of SEQ ID NO: 10 which forms a complex with HLA-C clone 10 molecules, comprising contacting a sample from a subject with an agent specific for said complex and determining interaction between said complex and said agent as a determination of said disorder.
28. Method for diagnosing a disorder characterized by expression of a tumor rejection antigen precursor coded for by a nucleic acid molecule having the sequence set

5

forth in SEQ ID NO: 1, comprising contacting a sample from a subject with an agent specific for a tumor rejection antigen derived from said precursor and consisting of the amino acid sequence of SEQ ID NO: 10, and determining interaction between said agent and said sequence or said expression product as a determination of said disorder.

1/4











FIG. 2**FIG. 4**

FIG. 3



4/4

FIG. 5



INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/06534

A. CLASSIFICATION OF SUBJECT MATTER IPC(5) : Please See Extra Sheet. US CL : Please See Extra Sheet. According to International Patent Classification (IPC) or to both national classification and IPC				
B. FIELDS SEARCHED Minimum documentation searched (classification system followed by classification symbols) U.S. : 424/88; 435/7.0, 7.1, 69.3, 93V, 320; 530/326, 328, 350, 828; 536/27; 935/12 Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched Electronic data base consulted during the international search (name of data base and, where practicable, search terms used) APS, BIOTECH, BIOSIS, CA, DERWENT, EMBASE, DERWENT, MEDLINE, REGISTRY, WPIDS, search terms: Bage, Trap, Van der Bruggen, Boon, Coulie, Renault				
C. DOCUMENTS CONSIDERED TO BE RELEVANT				
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.		
X	PROGRESS IN IMMUNOLGY, Volume 8, issued 1992, T. Boon et al., "Genes Coding for Tumor Antigens", pages 871-876, see entire document.	1, 3, 4, 6		
X	IMMUNOGENETICS, Volume 35, issued 1992, C. Traversari et al., "Transfection and Expression of a Gene Coding for a Human Melanoma Anitgen Recognized by Autologous Cytolytic T Lyphocytes", pages 145-152, see entire document.	1 and 4		
P, A	SCIENCE, Volume 262, issued 05 November 1993, J. Cohen, "Cancer Vaccines Get a Shot in the Arm", pages 841-843, see entire document.	15-18, and 22-26		
<input checked="" type="checkbox"/> Further documents are listed in the continuation of Box C. <input type="checkbox"/> See patent family annex.				
<table border="0"> <tr> <td> * Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier documents published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed </td> <td> *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *A* document member of the same patent family </td> </tr> </table>			* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier documents published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *A* document member of the same patent family
* Special categories of cited documents: *A* document defining the general state of the art which is not considered to be of particular relevance *E* earlier documents published on or after the international filing date *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) *O* document referring to an oral disclosure, use, exhibition or other means *P* document published prior to the international filing date but later than the priority date claimed	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art *A* document member of the same patent family			
Date of the actual completion of the international search 08 AUGUST 1994		Date of mailing of the international search report SEP 02 1994		
Name and mailing address of the ISA/US Commissioner of Patents and Trademarks Box PCT Washington, D.C. 20231 Facsimile No. (703) 305-3230		Authorized officer ANTHONY C. CAPUTA <i>Joe Warden for</i> Telephone No. (703) 308-0196		

INTERNATIONAL SEARCH REPORTInternati nal application No.
PCT/US94/06534

C (Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	SCIENCE, Volume 254, issued 13 December 1991, P. Van Der Bruggen et al., "A Gene Encoding an Antigen Recognized by Cytolytic T. Lymphocytes on a Human Melanoma", pages 1643-1647, see entire document.	1, 3, 4, 6

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US94/06534

A. CLASSIFICATION OF SUBJECT MATTER:
IPC (5):

A61K 37/02; C07H 21/04; C07K 7/06, 7/08, 15/06; C12N 15/03, 15/06; C12P 21/02; C12Q 1/68

A. CLASSIFICATION OF SUBJECT MATTER:
US CL :

424/88, 93V; 435/6, 7.1, 69.3, 320; 530/326, 328, 350, 828; 536/27; 935/12